

Spin-dependent lifetimes of hot electrons in 3d ferromagnets

K. M. Döbrich¹, A. Goris^{1,2}, I. Panzer¹, A. B. Schmidt^{1,3},
M. Pickel^{1,3}, M. Donath³, and M. Weinelt^{1,2}

¹*Max-Born-Institut, Max-Born-Straße 2A, 12489 Berlin, Germany*

²*Institut für Experimentalphysik, Freie Universität Berlin,
Arnimallee 14, 14195 Berlin, Germany*

³*Physikalisches Institut, Westfälische Wilhelms Universität,
Wilhelm-Klemm-Straße 10, 48149 Münster, Germany*

The spin-dependent lifetimes of electrons excited to energies of a few hundred meV above the Fermi energy E_F —so-called hot electrons—of the 3d ferromagnets Fe, Co and Ni have been investigated experimentally and theoretically for more than a decade. They are of high technological relevance through the associated giant magneto-resistance effect and are of fundamental interest—some of the underlying microscopic physical processes are not yet fully understood. The first and up to now only spin-resolved two-photon photoemission (SR-2PPE) measurements have been performed in Refs. [1,2], where thin metal films were grown on Cu(100) and the surfaces were covered with Cs to lower the work functions.

We present bichromatic infrared pump and ultraviolet probe SR-2PPE data that give experimental access to smaller energies above E_F and thereby extend the experimental data range, without the need of Cs adsorption. We find good agreement with the previously measured data [1,2] in the overlapping energy range and discuss discrepancies with recent *ab-initio* calculations. A possible reason for the discrepancies are spin-flip exchange-scattering processes that we could observe on the clean, i.e. Cs-free, Co surface.

[1] M. Aeschlimann, M. Bauer, S. Pawlik, W. Weber, R. Burgermeister, D. Oberli, and H. C. Siegmann, Phys. Rev. Lett. **79**, 5158 (1997).

[2] R. Knorren, K. H. Bennemann, R. Burgermeister, and M. Aeschlimann, Phys. Rev. B **61**, 9427 (2000).