

Ultrafast, Element-Specific, Demagnetization Dynamics Probed using Coherent High Harmonic Beams

S. Mathias^{1,2}, C. La-O-Vorakiat¹, P. Grychtol³, R. Adam³, M. Siemens¹, J. M. Shaw⁴, H. Nembach⁴, M. Aeschlimann², C. M. Schneider³, T. Silva⁴, M. Murnane¹, H. C. Kapteyn¹,

¹ *JILA, University of Colorado and NIST, Boulder, Co, USA*

² *Research Center OPTIMAS, University of Kaiserslautern, Germany*

³ *Institute of Solid State Research, IFF-9, FZ Jülich, Germany*

⁴ *Electromagnetics Division, NIST, Boulder, Co, USA*

Element-specific magnetization dynamics on nanometer length scales and femtosecond timescales is a topic of intense current interest. Questions on, e.g., femtosecond magnetization processes, ultrafast dynamics of nanodomains, or ultrafast interaction in magnetic multilayer systems have motivated a variety of studies using magneto-optic effects probed by ultrafast light pulses. Magneto-optical dynamics studies currently make use of either ultrafast lasers or x-rays from large-scale synchrotrons. Ultrafast pulses from femtosecond lasers (~ 20 fs) make high time resolution easily accessible, however, these studies lack element selectivity. X-rays from synchrotrons, on the other hand, enable element-specific probes using photons with energy near the absorption edges of ferromagnetic materials. Moreover, nanometer imaging capabilities are also possible using x-rays. However, the relatively poor time resolution available from x-ray facilities (~ 100 fs using femtosecond slicing) to date has been too slow to resolve the fastest magnetization dynamics.

Ultrafast, coherent, table-top, x-ray sources based on high-harmonic upconversion of femtosecond lasers promise to overcome these limitations by providing a new tool to study how magnets work at the shortest time and length scales, with element specificity. In this work, we use few-femtosecond extreme ultraviolet (EUV) pulses from high harmonic generation (HHG) to extract demagnetization dynamics and hysteresis loops of a compound material for the first time [La-O-Vorakiat *et al.*, PRL 103, 257402 (2009)]. We measure the fastest, elementally-specific, demagnetization dynamics of any approach to date, with a time resolution of 55 fs.




Figure 1: Time-resolved measurement of ultrafast demagnetization in a Permalloy film probed using few-fs high-harmonic beams around the Ni (blue circle) and Fe (red square) M-edge.