

# Electronic properties of graphene on Ir(111)

P. Pervan<sup>1</sup>, I. Pletikosić<sup>1</sup>, M. Kralj<sup>1</sup> M. Milun<sup>1</sup>

<sup>1</sup>*Institute of Physics, Bijenička 46, HR-10000 Zagreb, Croatia*

Epitaxial graphene on Ir(111) prepared in excellent structural quality[1] appears to be an attractive model system for the investigation of intrinsic electronic properties of graphene modified by weak interactions with a supporting surface[2]. Angle resolved photoemission spectroscopy (ARPES) clearly displays a Dirac cone, a hallmark of the unusual band structure of graphene[3], with the Dirac point shifted only slightly above the Fermi level which indicates marginal substrate induced doping. The Moiré structure resulting from the overlaid graphene and Ir(111) surface lattices imposes a superperiodic potential giving rise to opening of minigaps in the band structure. ARPES of the electron doped graphene reveals a whole variety of many body interactions[4]. The  $\pi^*$  band of potassium intercalated graphene showed noticeable renormalization within the wide energy range. Near the Fermi level the renormalization is induced by electron-phonon interactions while the change of dispersion at higher binding energies is induced by the hole-plasmon interaction.

[1] J Coraux, AT N'Diaye, M Engler, C Busse, D Wall, N Buckanie, FJM zu Heringdorf, R van Gastel, B Poelsema and T Michely, *New Journal of Physics* **11** 023006 (2009)

[2] I Pletikosić, M Kralj, P Pervan, R Brako, J Coraux, A N'Diaye, C Busse and T Michely *Phys. Rev. Lett.* **102** 056808 (2009)

[3] KS Novoselov, AK Geim, SV Morozov, D Jiang, MI Katsnelson, IV Grigorieva, SV Dubonos and A Firsov *Nature* **438** 197 (2005)

[4] A Bostwick, H Ohta, T Seyller, K Horn, O Rotenberg, *Nature Physics* **3** 35 (2007)